

---

---

## Spatial Distribution of Seafloor Biogeological and Geochemical Processes as Proxy to Evaluate Fluid-Flux Regime and Time Evolution of a Complex Carbonate/Hydrates Mound, Northern Gulf of Mexico

Leonardo Macelloni<sup>1</sup>, Simona Caruso<sup>2</sup>, Laura Lapham<sup>3</sup>,  
Carol B. Lutken<sup>1</sup>, Charlotte Brunner<sup>4</sup>, and Allen Lowrie<sup>5</sup>

<sup>1</sup>Mississippi Mineral Resources Institute, Center for Marine Resources and Environmental Technology and Seabed Technology Research Center, University of Mississippi, 310 Lester Hall, University, Mississippi 38677

<sup>2</sup>Fugro Survey Limited, Survey House, Denmore Road, Bridge of Don, Aberdeen, Scotland

<sup>3</sup>Center for Geomicrobiology, Aarhus University, Ny Munkegade 114-116, Bldg. 1540, DK-8000 Aarhus C, Denmark

<sup>4</sup>Department of Marine Sciences, University of Southern Mississippi, 1020 Balch Blvd., Stennis Space Center, Mississippi 39529

<sup>5</sup>238 F. Z. Goss Rd., Picayune, Mississippi 39466

---

---

### ABSTRACT

A one km diameter carbonate/hydrate mound in Mississippi Canyon Block 118 is the site of the Gulf of Mexico Hydrates Research Consortium (GOMHRC) multi-sensor, multi-discipline seafloor observatory (SFO). In preparation for installing the observatory and understanding mound physiography and dynamics, geophysical, biological, geological and geochemical surveys have been carried out. By integrating high resolution swath bathymetry, acoustic imagery, seafloor video, and shallow geological samples in a morpho-biogeological model we have described a complex mound structure consisting of three main crater clusters: northwest, southwest, and southeast, each associated with a distinct, deep-seated fault. The crater complexes exhibit differences in morphology, bathymetric relief, outcropping hydrates, venting fluids, seafloor sediments, and biological community patterns. Spatial distribution of these attributes suggests that the craters represent three different stages of fluid flux regime; the southeast crater seems to be an extinct or quiescent vent, the southwest a mature, fully open vent and the northwest a recently established vent. Geochemical evidence from pore-water gradients corroborates this model suggesting that upward fluid flux waxes and wanes over time and that microbial activity is sensitive to such change. Sulfate and methane concentrations show that microbial activity is patchy in distribution and is typically higher within the northwest and southwest complexes, and diminished significantly over the southeast complex. Three distinct fluid-flux regimes coexisting within a single mound complex confirm the necessity of continuous monitoring over the five-to-ten years that the observatory is expected to be operational.